

Modified Nicoll's Graft for Treatment of Gap Non-union of Ulna: A Rare Case Report

Chaitanya Krishna¹, Jayanta Mondal¹, Anirudhha Sengupta¹

What to Learn from this Article?

Nicoll's graft is ideal for gap non-union of bones like radius and ulna where defect is up to 3 cm.

Abstract

Introduction: Despite the multitude of studies for the treatment of non-unions, no clear criterion has been established for treating a fracture of non-union. Non-union of radius and ulna produces a discrepancy in length which must be restored for adequate rotation and function. Nicoll described a technique of bridging gaps in long bones with solid blocks of cancellous bone and fixing the fragments with metal plates. We utilized modified Nicoll's graft using corticocancellous graft taken from iliac crest.

Case Report: In the present case, a 17-year-old male patient of eastern India presented with gap non-union of 3 cm in ulna of non-dominant hand, left side after 3 years and 6 months of trauma. There was linear surgical scar mark of 8 cm overlying ulna, but there was no sign of infection. He was functionally crippled with severe restriction of range of motion.

Conclusion: The tricortical cancellous bone graft under optimal compression provides a good solution for the management of gap non-union of both bone forearm for deficit up to 3 cm with no evidence of infection.

Keywords: Both bone, gap non-union, Nicoll's graft, ulna.

Introduction

Orthopedic surgeons can choose from numerous surgical and nonsurgical methods that vary greatly in their invasiveness and propensity for minor and catastrophic complications in the treatment of gap non-union. In a situation with equal risks and benefits, the simplest, most easily tolerated method should be chosen. For many years, the most frequently used method of treatment of non-unions has been bone grafting. Autogenous cancellous bone, although limited in quantity and associated with significant donor site morbidity, remains the "gold standard" in grafting material.

Various treatment modalities described in treating forearm bone loss include using boiled cadaveric bones [1], use of cortical tibial graft

with screws [2], grafts using ulnar segment held by screws [3], use of iliac crest graft to fill the bone gap and fixation with an intramedullary nail [4], cancellous insert grafts with plate fixation [5, 6], bone transport in forearm bones using the principles of ilizarov [7], using fibula as an intercalary bone graft and with a tibial cortical bone graft fixed opposite to a plate [5, 8], and vascularized fibular iliac bone graft [9].

In 1956, Nicoll [5] described a method of overcoming this problem by filling the defect left by excision of dead bone with cancellous bone from the anterior iliac crest, securing the graft by a plate and screws. In 1980, grace and Evers modified Nicoll's technique using a corticocancellous graft, fixed by a plate.

Author's Photo Gallery



Dr. Chaitanya Krishna



Dr. Jayanta Mondal



Dr. Anirudhha Sengupta

Access this article online

Website:
www.jocr.co.in

DOI:
2250-0685.730

¹Department of Orthopedics, Medical College and Hospital, College Street, Kolkata, West Bengal, India.

Address of Correspondence

Dr. Chaitanya Krishna,
Fellow in Joint Replacement and Reconstruction Centre, Sir Ganga Ram Hospital, New Delhi, India.
E-mail: chatconfident@gmail.com

The present paper discusses results of gap non-union of ulna treated by tricortico-cancellous bone graft with plate fixation.

Case Report

A 17-year-old male patient came to us with severe restriction of movement of left elbow joint and inability to do his daily activities. On examination, linear surgical scar mark was present over the left forearm on dorsal aspect; there was no sign of infection. The range of motion was 70-90° with restricted supination and pronation. On palpation, the metal plate was palpated over dorsal ulna. X-ray showed distinct gap non-union of ulna. There was no distal neurovascular deficit.

The patient had a history of road traffic accident 4 years back with fracture of both bone forearm proximal third (Monteggia variant) (Fig. 1). This fracture was internally fixed with narrow dynamic compression plate (DCP), but after 3 months implant breakage occurred of ulna (Fig. 2). Then, plate was removed and fixed with Rush nail (Fig. 3). After

10 months, there was no sign of union, so rush nail was removed and fixed with narrow DCP (Fig. 4). Even after 2 years, there was no sign of union as ends of bone were sclerotic with a distinct gap in between (Fig. 5).

Hence, we decided to go for modified Nicoll's graft. We put a corticocancellous strut graft taken from anterior iliac crest and put it at the gap site after freshening the ends and under compression with arthrolysis of the joint (Fig. 6). After 4 months, there was a union at fracture site with 20-120° range of movement and 50° supination and 40° pronation (Fig. 7). The patient now can do his daily activity and is very much satisfied (Fig. 8 and 9).

Discussion

A gap non-union is often characterized by significant scarring, sclerosis, and absorption of bone at fracture ends, leading to devascularization and lack of osteogenic potential due to biological failure. There are numerous




Figure 1: Radiograph of patient 4 years back with fracture both bone forearm proximal third.



Figure 2: Radiograph after 3 months with implant breakage of ulna.




Figure 3: Radiograph of a patient with plate removal and fixed with Rush nail of ulna.



Figure 4: Radiograph after 10 months with no sign of union, so Rush Nail was removed and fixed with narrow dynamic compression plate.




Figure 5: Even after 2 years, there was no sign of union as ends of bone were sclerotic with distinct gap in between.




Figure 6: Radiograph of the patient after putting Nicoll's graft.



Figure 7: Radiograph of the patient with the union of fracture after 4 months.




Figure 8: Radiograph of the patient in extension.



Figure 9: Radiograph of the patient in flexion.

treatment modalities of non-unions, but there is a paucity of the literature toward the management of atrophic non-unions of forearm bones with gaps.

Nicoll described a method to deal with the problem of gap non-union, by filling the defect left after the excision of the dead bone, with cancellous bone from the anterior iliac crest and fixing the graft by compression plate. There are very few published reports on Nicoll's technique. This simple technique has not gained its popularity and is not being practiced by many as a standard orthopedic procedure for the management of gap non-unions of single or both bones of the forearm.

The present study was undertaken to treat non-unions with a gap in forearm bones by modified Nicoll's technique. The clinical and experimental evidence has proved that a fresh autogenous cancellous bone graft is always better than any other modality of treatment. For physical reasons, corticocancellous graft gives a more rapid union and less fibroblastic reactions than chips or slices [5, 10]. The sclerotic ends of bone are devoid of blood supply and lose the osteogenic potential. Siffert [11] regarded the freshened bone ends as the most important single source of osteogenesis.

This technique of grafting is simple and provides stability to fracture as well as bone graft. The presence of cortical bones on three sides provides adequate strength with optimal compression at host-graft junction until the cancellous bone is involved in the formation of new bone due to creeping substitution and also allows for the early invasion of the periosteal circulation [12]. The abundance of the cancellous bone decreases the time of graft incorporation and filling up the gaps in the process of union. Thus, the procedure serves in

three ways, i.e., internal fixation, defect bridging, and osteogenesis promoter.

Conclusion

The Nicoll's bone graft which if modified with tricortico cancellous strut bone graft under optimal compression with the help of compression plate provides a high union rate for difficult problem of atrophic gap non-union of both bone forearms.

Clinical Message

Gap non-union can be challenging to treat. Many options are available with their own complications and prognosis. Nicoll's graft provides an excellent option with minimal intervention and early union. With almost 3 cm of gap non-union, this case report provides an excellent example of the use of Nicoll's graft.

References

- Lloyd-Roberts GC. Experiences with boilded cadaveric bone. *J Bone Joint Surg Br* 1952;34-B(3):428-432.
- Boyd HB. The Treatment of difficult and unusal non-union. *J Bone Joint Surg* 1943;25:535
- Miller RC, Phalen GS. The repair of defects of the radius with fibular bone grafts. *J Bone Joint Surg Am* 1947;29(3):629-636.
- Spira E. Bridging of bone defects in the forearm with iliac graft combined with intramedullary nailing. *J Bone Joint Surg Br* 1954;36-B(4):642-646.
- Nicoll EA. The treatment of gaps in long bones by cancellous insert grafts. *J Bone Joint Surg Br* 1956;38-B(1):70-82.
- Dabezies EJ, Stewart WE, Goodman FG, Deffer PA. Management of segmental defects of the radius and ulna. *J Trauma* 1971;11(9):778-788.
- Tetsworth K, Krome J, Paley D. Lengthening and deformity correction of the upper extremity by the Ilizarov technique. *Orthop Clin North Am* 1991;22(4):689-713.
- Moroni A, Rollo G, Guzzardella M, Zinghi G. Surgical treatment of isolated forearm non-union with segmental bone loss. *Injury* 1997;28(8):497-504.
- Jupiter JB, Gerhard HJ, Guerrero J, Nunley JA, Levin LS. Treatment of segmental defects of the radius with use of the vascularized osteoseptocutaneous fibular autogenous graft. *J Bone Joint Surg Am* 1997;79(4):542-550.
- Abbott LC, Schottstaedt ER, Saunders JB, De CM, Bost FC. The evaluation of cortical and cancellous bone as grafting material; A clinical and experimental study. *J Bone Joint Surg Am* 1947;29(2):381-414.
- Siffert RS. Experimental bone transplants. *J Bone Joint Surg Am* 1955;37-A(4):742-758.
- Gomez EA. Treatment of forearm non union with iliac graft and a Hunec nail. *Acta Ortop Maxicana* 2005;19:S28-S33.

Conflict of Interest: Nil.
Source of Support: None

How to Cite this Article

Krishna C, Mondal J, Sengupta A. Modified Nicoll's Graft for Treatment of Gap Non-union of Ulna: A Rare Case Report. *Journal of Orthopaedic Case Reports* 2017 Mar-Apr;7(2):17-20.